No.



8300143

### THE UNIVERD STAYIES OF AMERICA

TO ALL TO WHOM THESE PRESENTS SHALL COME:

# Pioneer Hi-Bred International, Inc.

Cancreas, there has been presented to the

#### Secretary of Agriculture

AN APPLICATION REQUESTING A CERTIFICATE OF PROTECTION FOR AN ALLEGED NOVEL VARIETY OF SEXUALLY REPRODUCED PLANT, THE NAME AND DESCRIPTION OF WHICH ARE CONTAINED IN THE APPLICATION AND EXHIBITS, A COPY OF WHICH IS HEREUNTO ANNEXED AND MADE A PART HEREOF, AND THE VARIOUS REQUIREMENTS OF LAW IN SUCH CASES MADE AND PROVIDED HAVE BEEN COMPLIED WITH, AND THE TITLE THERETO IS, FROM THE RECORDS OF THE PLANT VARIETY PROTECTION OFFICE, IN THE APPLICANT(S) INDICATED IN THE SAID COPY, AND WHEREAS, UPON DUE EXAMINATION MADE, THE SAID APPLICANT(S) IS (ARE) ADJUDGED TO BE ENTITLED TO A CERTIFICATE OF PLANT ARIETY PROTECTION UNDER THE LAW.

NOW, THEREFORE, THIS CERTIFICATE OF PLANT VARIETY PROTECTION IS TO GRANT INTO THE SAID APPLICANT(S) AND THE SUCCESSORS, HEIRS OF ASSIGNS OF THE SAID APPLICANT(S) FOR THE TERM OF CIGHTEEN YEARS FROM THE DATE OF THIS GRANT, SUBJECT TO THE PAYMENT OF THE REQUIRED REES AND PERIODIC REPLENISHMENT OF VIABLE BASIC SEED OF THE VARIETY IN A PUBLIC, REPOSITORY AS PROVIDED BY LAW, THE RIGHT TO EXTEND OTHERS FROM SELLING THE VARIETY, OR OFFERING IT FOR SALE, OR REPRODUCING IT, DRING IT, OR EXPORTING IT, OR USING IT IN PRODUCING A HYBRID OR DIFFERENT THEREFROM, TO THE EXTENT PROVIDED BY THE PLANT VARIETY PROTECTION ACT 2, AS AMENDED, 7 U.S.C. 2321 ET SEQ.)

CORN

'G50'

In Lestimony Winercot, I have hereunto set my hand and caused the seal of the Plant Variety Protection Office to be affixed at the City of Washington this 21st day of Vecember in the year of our Lord one thousand nine

the year of our Lord one thousand nine hundred and eighty-four.

John R Block

Ken not Steran

Plant Variety Protection Office Agricultural Marketing Service

XXAXXXXXXXXX

U.S. DEPARTMENT OF AGRICULTU AGRICULTURAL MARKETING SERV LIVESTOCK, MEAT, GRAIN & SEED DIV	FORM APPROVED: OMB NO. 0581-0006 No certificate for plant variety protection			
APPLICATION FOR PLANT VARIETY PROTECTIONS ON reverse)	CTION CERTIFICATE	may be issued unless a completed appli- cation form has been received (6 U.S.C. 553).		
1. NAME OF APPLICANT(S)	2. TEMPORARY DESIGNATION	3. VARIETY NAME		
Pioneer Hi-Bred International, Inc.		G50		
4. ADDRESS (Street and No. or R.F.D. No., City, State, and Zip Code) Plant Breeding Division		FOR OFFICIAL USE ONLY PVPO NUMBER		
Department of Corn Breeding P.O. Box 85, Johnston, IA 50131-0085	515/270-3300	8300143		
6. GENUS AND SPECIES NAME Zea mays  7. FAMILY NAM Gramine		DATE 6/1/83 TIME 8:30 NA.M. P.M.		
8. KIND NAME 9.	DATE OF DETERMINATION	AMOUNT FOR FILING		
Corn	1978	DATE 6/1/83  AMOUNT FOR CERTIFICATE		
10. IF THE APPLICANT NAMED IS NOT A "PERSON," GIVE FORM partnership, association, etc.)  Corporation	OF ORGANIZATION (Corporation	\$ 500.00		
11. IF INCORPORATED, GIVE STATE OF INCORPORATION IOWA		11/15/84  12. DATE OF INCORPORATION May 6, 1926		
14. CHECK APPROPRIATE BOX FOR EACH ATTACHMENT SUBMIT  Exhibit A, Origin and Breeding History of the Variety (See Section 52 of the Plant Variety Protection Act.)  b. X Exhibit B, Novelty Statement	c. Exhibit C; Objective I from Plant Variety Pr	Description of the Variety (Request form otection Office.)  Description of the Variety		
15. DOES THE APPLICANT(S) SPECIFY THAT SEED OF THIS VARI SEED? (See Section 83(a) of the Plant Variety Protection Act.)		E ONLY AS A CLASS OF CERTIFIED items 16 and 17 below) X No		
16. DOES THE APPLICANT(S) SPECIFY THAT THIS VARIETY BE LIMITED AS TO NUMBER OF GENERATIONS?	17. IF "YES" TO ITEM 16,	WHICH CLASSES OF PRODUCTION ED?		
☐ Yes ☐ No	Foundation	Registered Cartifled		
18. DID THE APPLICANT(S) FILE FOR PROTECTION OF THE VAR	IETY IN THE U.S. OR OTHER CO	Yes (If "Yes," give name of countries and dates)		
		X No		
19. HAVE RIGHTS BEEN GRANTED IN THE U.S. OR OTHER COUN	TRIES?	Yes (If "Yes," give name of countries and dates)		
		X No		
20. The applicant(s) declare(s) that a viable sample of basic seed plenished upon request in accordance with such regulations:	s of this variety will be furnished as may be applicable.	d with the application and will be re-		
The undersigned applicant(s) is (are) the owner(s) of this sex distinct, uniform, and stable as required in Section 41, and is Variety Protection Act.	ually reproduced novel plant va	riety, and believe(s) that the variety is ne provisions of Section 42 of the Plant		
Applicant(s) is (are) informed that false representation herei	n can jeopardize protection and	result in penalties.		
Pioneer Hi-Bred International, Inc.		DATE		
by: SIGNATURE OF APPLICANT		DATE		
Richard & Melannell		May 26, 1983 1		

'G50'

14A. Exhibit A. Origin and Breeding History

Pedigree: 848/207)X1321X

Pioneer line 'G50', Zea mays L., a yellow dent corn inbred, was developed by Pioneer Hi-Bred International, Inc. from the F2 population of the single cross  $848 \times 207$ . The progenitors of 'G50' are also proprietary inbred lines of Pioneer Hi-Bred International, Inc. The pedigree method of breeding was used in the development of this inbred as per the following.

F2 seed was obtained in the field at Tipton, Indiana, during the summer of 1974 by selfing the F1 hybrid,  $848 \times 207$ . population was grown and self-pollinated at Algona, Iowa, in 1975. Twenty-three ears were saved from the F2 population. This F3 seed was grown ear-to-row at Homestead, Florida, during the winter of 1975-76. Five self-pollinated ears were saved from F3 ear-row number one. The F4 family was grown ear-to-row at Algona in 1976, and two self-pollinated ears from ear-row number three were saved. During 1977, the F5 generation was grown at Algona, and two selfpollinated ears were saved from the F5 ear-row number two. addition, the F5 was crossed to an inbred tester for the purpose of getting an estimate of the line's general combining ability. In 1978, yield trials were conducted at Algona, Iowa, involving the testcross made in 1977 to the F5. In the nursery, three ears were saved from ear-row number one from the F5 generation. Additional crosses were made for yield testing in 1979. Based on hybrid yield test performance and the line's per se performance in the nursery, it was determined that this line was superior to other inbreds evaluated in 1978, and it was named 'G50' December 1978. Since the time that the line was named, additional hybrid combinations have been evaluated and subsequent generations of the line have been grown and hand-pollinated with observations made for uniformity. An outline of the breeding profile of the inbred is attached.

'G50' has shown uniformity and stability for all traits as described in Exhibit C (form LPGS-470-28) - "Objective Description of Variety." It has been self-pollinated and ear-rowed a sufficient number of generations with careful attention paid to uniformity of plant type to assure genetic homozygosity and phenotypic stability. 'G50' has been increased by the Parent Corn Department, Pioneer's foundation seed group, each year since 1981. The line has been increased both by hand and in isolated fields with continued observation for uniformity.

Page 2

#### 14A. Exhibit A. Origin and Breeding History (continued)

Two variant traits have been observed in 'G50'. Under certain environmental conditions at maturity, the pericarp tends to "pop" which predisposes the ear to molds. In addition, it has been noticed in current Parent Corn seed lots that between 25 to 35 percent of the kernels lack the usual complement of embryo leaves. These abnormal kernels germinate, but the resultant seedlings die soon after emergence. This abnormality appears to be due to recessive gene action. Breeding techniques are currently being undertaken to identify an ear-row segregate that does not carry this variant trait.

Pioneer Hi-Bred International, Inc., Des Moines, Iowa, is the employer of the plant breeders involved in the selection and development of  $^{\prime}G50^{\prime}$ . Pioneer Hi-Bred International, Inc. has the sole rights and ownership of  $^{\prime}G50^{\prime}$ .

14A. Exhibit A. Origin and Breeding History of Corn Inbred Line 'G50'

Season/Year	Inbreeding Level	Nursery Location	Pedigree	Ears Saved
Summer 1973	F0	Tipton, IN	Fl cross made.	Bulk
Summer 1974	Fl	Tipton, IN	848/207	Bulk
Summer 1975	F2	Algona, IA	848/207)X	23
Winter 1975-76	F3	Homestead, FL	848/207)X1	5
Summer 1976	F4	Algona, IA	848/207)X13	2
Summer 1977	F5*	Algona, IA	848/207) X132	.2
Summer 1978	F6**	Algona, IA	848/207)X1321	3
December 1978	Line nam	ed 'G50'.	• •	
1978-1982		fields with obs	pollination and in servations made for	

<sup>\*</sup>Testcross made for yield testing in 1978.

<sup>\*\*</sup>More hybrid combinations made involving 'G50' for testing in 1979.

### 4B. Exhibit B. Novelty Statement

'G50' is most similar to the inbred line 207. 207 is also a proprietary inbred line of Pioneer Hi-Bred International, Inc. and is half the parentage of 'G50'. 'G50' differs from 207 in maturity, glume color, and silk color. 'G50' reaches 50% pollen shed and 50% silk, 115 and 100 heat units, respectively, later than 207. 'G50's' glume color is green whereas the glume color for 207 is red. The silk color of 'G50' is green; the silk color of 207 is red.

## OBJECTIVE DESCRIPTION OF VARIETY CORN (ZEA MAYS)

NAME OF APPLICANTIS	FOR OFFICIAL USE ONLY
Pioneer Hi-Bred International, Inc.	P VPO NUMBER
ADDRESS (Street and No. or R.F.D. No., City, State, and ZIP Code)	8300143
Plant Breeding Division	VARIETY NAME OR TEMPORARY DESIGNATION
Department of Corn Breeding PO Box 85	'G50'
Johnston, IA 50131-0085	· · · · · · · · · · · · · · · · · · ·
Place the appropriate number that describes the varietal character of this variety in the Place a zero in first box (e-s- $089$ or $9$ or $9$ ) when number is either 99 or less or	
1. TYPE:	
2 1 = SWEET 2 = DENT 3 = FLINT 4 = FLOUR 5 = PC	OP 6 = ORNAMENTAL
2. REGION WHERE BEST ADAPTED IN THE U.S.A.:	
1 = NORTHWEST 2 = NORTHCENTRAL 3 = NORTHEAST 5 = SOUTHCENTRAL 6 = SOUTHWEST 7 = MOST REGIONS	4 = SOUTHEAST
	comments" (pg. 3) state how
	s were calculated)
6 6 DAYS FROM EMERGENCE TO 50% OF PLANTS IN SILK	9 3 HEAT UNITS
DAYS FROM 50% SILK TO OPTIMUM EDIBLE QUALITY	HEAT UNITS
DAYS FROM 50% SILK TO HARVEST AT 25% KERNEL MOISTURE	HEAT UNITS
4. PLANT:	
2 2 5 CM. HEIGHT (To tassel tip)	0 0 CM, EAR HEIGHT (To base of top
0 8 CM. LENGTH OF TOP EAR INTERNODE	
Number of Tillers: Number of Ears Per Stalk:	:
	SLIGHT TWO-EAR TENDENCY
1 1 1 1 NONE 2 1 1 2 3 2 2 3 4 3 1 4 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	-EAR TENDENCY 4 = THREE-EAR TENDE
Cytoplasm Type:	
1 1 = NORMAL 2 = "T" 3 = "S" 4 = "C" 5 = OTHER	(Specify)
5. LEAF (Field Corn Inbred Examples Given):	
Calcu	
Observed olive green  1 = Light Green (HY) 2 = MEDIUM GREEN (WF9) 3 = DARK GREEN	EEN (B14) 4 = VERY DARK GREEN (
Angle from Stalk (Upper half):  Sheath Pubscence:	
1 = $<30^{\circ}$ 2 = $30-60^{\circ}$ 3 = $>60^{\circ}$ 1 = LIGHT 3 = HEAVY	
Marginal Waves: Longitudinal Creases:	
2 1 = NONE (HY) 2 = FEW (WF9) 3 = MANY (OH7L) 1 1 = ABSEN 3 = MANY (	•
Width: Length:	
0 9 CM. WIDEST POINT OF EAR NODE LEAF 0 8 6 CM. E.	AR NODE LEAF
2 0 NUMBER OF LEAVES PER MATURE PLANT	<i>₽</i>

			· · · · · · · · · · · · · · · · · · ·			· · · · · · · · · · · · · · · · · · ·			43/1/1 <del>1</del> /
6.	TASSEL:	•	-		-		4.4	•	
- - -	1 1	NUMBER OF I	LATERAL BRANC	CHES			* **		
					4	*			
	Branch Ang	ie from Central S	pike:	e a company of the co	Pendur	ncle Length:	•		
	2	1 = < 30?	2 = 30 40°	3 = > 45°		1 1	CM. FROM TO	P LEAF TO BASAL	BRANCHES
	Polien Shed	l:							
-	2	1 = LIGHT (WI	Ea)	2 = MEDIUM		3 = HEAVY	(KY21)		
	ت								
			)	lsh-purple,	seconda	ry oran	ge-yellow		
	<u> </u>	Anther Color:	7			3 = REC	4=1	PURPLE	= GREEN
	5	Observed	6 = OTHER ( l olive gree	Specify)			<del></del>	· · · · · · · · · · · · · · · · · · ·	
	Pollen Rest			rted, 1 = Partial, 2 =	Good)				•
	0т	0	"s" 0	"c"	OTHER (	Specify Cyto	piasm and degre	es of restoration)	
	•				e				
<del></del> 7	. EAR (Hus	ked Ear Data Exc	ept When Stated O	therwise):					
		· · · · · · · · · · · · · · · · · · ·							
	1 6	CM LENGTH		I, MID-POINT AMETER		9 8	GM. WEIGHT		
	Kernel Row	<b>!s</b> :					•		
	2	-		5.5555		1 4	NUMBER		
		1 = INDISTING	<u> 2</u> =	DISTINCT					
					- ** 	· · ·			
	2	1 = STRAIGHT	r 2.≂SL	IGHTLY CURVED	•	3 = SPIRA	L.		
	Silk Color (	Exposed at Silkin	ig Stage):				•		
		4 CDEEN	o – DINK	7 CAL	MON	4 = F			
	لئا	1 = GREEN Observed	2 = PINK greenish-ye	3 = \$AL	WION .	4-6	IED .		
	Husk Color	*	,						
		FRESH )	1 = LIGH	T GREEN	2 =	DARK GRI	EEN	3 = PINK	
	٢٢		}						
•	6	DRY	4 = RED	5	- PURPLE		6 = BUFF	4	•
	Husk Extent	tion: (Hervest St	age)		Husk L	eaf:			
		HORT (Ears Expo		A (Sarety Covering I	Ear)		= SHORT ( < 8 5 = LONG ( > 15	<b>.</b>	M (8-15 CM)
	4 = V	ERY LONG (>1			. 7.			, <b>G</b> , , ,	
	Shank:				PUSITIO	n at Dry Hu:	sk Stage:		6. Company of the Com
	1 0	CM LONG	8 NO. OF IN	TERNODES		1 1	= UPRIGHT	2 = HORIZONTA	L 3 = PENDENT
	Taper:				Drying	Time (Unho	ısked Ear): ्		
	2	· · · · · · · · · · · · · · · · · · ·			_ ,3			·	
	<u> </u>	1 = SLIGHT	2 = AVERAGE	3 = EXTREME	······································		= SLOW	2 = AVERAGE	3 = FAST
8.	KERNEL (C	•							•
•	aise (L.com)	Ear Mid-Point):			[				
		MM LONG	0 8	MM. WIDE	0 5	мм. тн	ICK		
	Shape Grade	(% Rounds)					•		
	3	1 = < 20	2 = 2040	3 = 40-6	50	4 = 60 -6	30	5 = > 80	

8. KERN	EL (Dried) :		_	0000143
1				B = TAN 4 = BRONZE 7 = CHERRY RED
	8 = V	ARIEGATED (Describe)		
				<del></del>
1	Aleurone Color: 1 = H	OMOZYGOUS 2 =	SEGREGATING (Describe)	
· —				
	1 = WHITE 2	= PINK 3 = TAN	4 = BROWN	<b>5 = BRONZE</b> 6 = RED
لتا	7 = PURPLE 8	= PALE PURPLE 9 =	VARIEGATED (Describe)	
3	Endosperm Color: 1	- WHITE 2 - PALE YELL	OW 3 - YELLOW	4 = PINK-ORANGE 5 = WHITE CAP.
	Observed orang	ge-yellow		
Endos	perm Type:	·		
	1 = SWEET (su 1)	2 = EXTRA SWEET (sh2)	3 = NORMAL STAR	CH 4 = HIGH AMYLOSE STARCH
3	5 = WAXY STARCH	6 = HIGH PROTEIN	7 = HIGH LYSINE	8 = OTHER (Specify)
2 3	7			
	GM. WEIGHT /100 SEED	S (Unsized Sample)	•	
9. COB:				
2 2	MM. DIAMETER AT MID	P-POINT		
Strengt	l ⊁h ·		Color:	
21,9118	1 = WEAK 2 = 5	TRONG		PINK 3 = RED 4 = BROWN
2	I-WEAR 2-5	INDING		6 OTHER (Specify) Reddish-orang
10				O C. T.C. Topoch y
IU. DISEA	SE RESISTANCE (O = Not 1	Tested, 1 = Susceptible, 2 = Resident Tole 1		
2	STALK ROT (Diplodia)	1 21	OT (Fuserium)	2 STALK ROT (Gibberella)
	STACK NOT (DIDIOGIE)		O' (Possilum)	
	NORTHERN LEAF BLIG	HT L SOUTHER	RN LEAF BLIGHT	2 smut (Head)
0	SOUTHERN RUST	2 CORN SM	υτ (Common)	2 BACTERIAL WILT (Stewart's)
	BACTERIAL LEAF BLIG	HT 1 MAIZE D	WARF MOSAIC	0 STUNT
٥	DACITAL CEAP BLIC	1	MARE MUSAIC	U STON!
	OTHER (Specify)		•	
11. INSEC	T RESISTANCT (O = Not Te	sted, 1 = Susceptible, 2 = Besists	unt):	
		Toleran		
2				<del></del>
يكا	cornsorer (European)	0 EARWORM	0 SAPBE	ETLE 0 APHID
0	ROOTWORM (Northern)	1 ROOTWORM (Weste	rn)	
		<b>=</b>	,	
0	ROOTWORM (Southern)	OTHER (Specify)	<u> </u>	
12. VARIE	TIES MOST CLOSELY RESE	MBLING THAT SUBMITTED I	FOR THE CHARACTERS GIV	FN-
CHARA		VARIETY	CHARACTER	VARIETY
Maturit		A632		
Plant T	<u> </u>	207	Kernel Type	207
Ear-Typ		207	Quality (Edible)	207
- CO1-54 AT	76		Usage	
REFEA	ENCES:			
	U.S. Department Agricultu	the state of the s		
	· ·	Products. 1970 Avi Publishing (		
		e, and A.C. Fraser. A Summary (	•	nell A.E.S., Mem. 180, 1935.
	•	58. Crop Science Society of Ame		
		red Lines of Ohio, Ohio A.E.S.		Ohio Coon Hair-d
·	<del></del>	stem for the Classification of Cor		
COMME	NTS: Heat units a	re accumulated from	daily temperature	s as follows: ot greater than 86.
	LO = Minimum	air temperature in	Februarie L.	
	Heat units =	$(HI + L\ddot{0})/2 - 50,$	but not less than	ot less than 50. ).
	and the second s	<b>_</b> _,		•

14D. Exhibit D. Additional Description of 'G50'

'G50' is a yellow dent inbred line of corn, Zea mays L.

As an inbred per se, 'G50' is similar to 207 in a number of plant and seed characteristics. Certain similarities are expected since half the parentage of 'G50' is 207. However, there are a number of distinguishable differences between the two inbreds as already stated in Exhibit B.

'G50'is similar to 207 in plant height and leaf color. 'G50' is, however, higher eared than 207 (80 centimeters versus 100 centimeters). 'G50' also has a leaf angle of less than 30 degrees whereas 207 has a leaf angle of between 30 and 60 degrees. 'G50' has about three more leaves per mature plant (20 versus 17 for 207. 'G50' has a shorter shank than 207 (10 centimeters versus 16 centimeters). 'G50' has a shorter tassel peduncle than 207 (11 centimeters to versus 22 centimeters). 'G50' also has harder textured kernels and a higher level of tolerance to European corn borer than 207. For comparitive purposes, data are attached with 'G50' and 207 crossed to the same tester inbred lines and evaluated in the same locations.

'G50' has average to above average tolerance to Helminthosporium leaf spot (Helminthosporium carbonum), eye spot (Kabatiella zeae), anthracnose (Colletotrichum graminicola), common rust (Puccinia sorghi), Stewart's bacterial wilt (Erwinia stewartii), Goss's wilt (Corynebacterium nebraskense), and head smut (Sphacelotheca reiliana). It has average to below average tolerance to Southern leaf blight (Helminthosporium maydis) and to the MDM virus complex. It has below average tolerance to Northern leaf blight (Helminthosporium turcicum), gray leaf spot (Cercospora zeae), sorghum downy mildew (Sclerospora sorghi), and corn lethal necrosis virus disease. 'G50' has above average tolerance to the first brood of European corn borer, but it is below average for tolerance to the second brood of European corn borer. It has a rather low level of tolerance to Lasso - Dual (alachlor - metolachior) herbicides.

'G50' makes high-eared hybrids that have stable yields under a wide range or environmental conditions. These hybrids are not only high yielding, but they dry down rapidly after reaching physiological maturity. 'G50' hybrids have average root quality and are above average for stalk quality. Grain of these hybrids is hard textured, of excellent quality, and high in test weight. Drought tolerance and late-season plant health of 'G50' crosses are above average.

Exhibit D. Comparison of 'G50' and 207 crossed to the same tester lines and the hybrids evaluated at the same locations. All values are expressed as percent of the test mean except yield, which is expressed as bushels/acre adjusted to 15.5% grain moisture (1980 data).

			% -	
Ear Height	39	104	66	9
Plant Height	39	100	66	1
Seedling Vigor	22	65	101	6
Cob Scores	21	119	108	11
Grain Quality	74	110	16	13
Test Weight	81	103	101	7
Stay Green	37	116	120	7
Fars/Plot	35	101	101	0
Root Lodging	99	104	112	8
Stalk Lodging	81	105	105	0
CDN SPeq	26	103	100	3
Moisture	93	66	98	-
Percent Yield	93	101	102	Ħ
Vield	93	134	136	2
Inbred		'G50'	207	
	No. of Reps.			Diff.

Exhibit D. Comparison of 'G50' and 207 crossed to the same tester lines and the hybrids evaluated at the same locations. All values are expressed as All values are expressed as is expressed as bushels/acre percent of the test mean except yield, which adjusted to 15.5% grain moisture (1981 data). 14D.


	·			
		,	:	:
Ear Height	174	103	101	2
Plant Height	174	86	100	2
Seedling Vigor	213	105	100	5
Cob Scores	118	101	104	3
Grain Quality	298	114	102	12
Test Weight	354	102	100	2
Stay Green	174	96	86	2
Ears/Plot	105	102	100	2
Root Lodging	144	101	104	3
Stalk Lodging	367	101	102	1
cDN Shed	65	100	66	1
Moisture	367	6	86	1
Percent Yield	367	102	102	0
Yield	367	142	142	0
Inbred		'G50'	207	
	No. of Reps.		-	Diff.

Comparison of 'G50' and 207 crossed to same tester lines and the hybrids evaluated at the same locations. All values are expressed as percent of the test mean except yield, which is expressed as bushels/acre adjusted to 15.5% grain moisture (1982 Data). Exhibit D. 14D.

				<u> </u>
		-		
Ear Height	85	102	66	4
Plant Height	85	100	100	0
Seedling Vigor	101	92	66	7
Cob Scores	38	94	103	6
Grain Quality	167	114	100	14
Test Weight	215	104	101	3
Stay Green	66	26	66	2
Fars/Plot	64	101	86	3
Root Lodging	107	102	103	<b>-</b> i
Stalk Lodging	215	101	103	2
cDu shed	39	100	66	-
Moisture	251	98	66	-
Percent Yield	215	101	100	
Vield	215	147	146	
pəıquI		'650'	207	
	No. of Reps.			Diff.

14D. Exhibit D. Additional Description of G50 (continued)

A. Whole plant



14D. Exhibit D. Additional Description of G50 (continued)

B. Tassel



14D. Exhibit D. Additional Description of G50 (continued)

C. Ear

